

DO QUARK MASSES RUN?

Arcadi SANTAMARIA

*Departament de Física Teòrica, IFIC, Universitat de València-CSIC, 46100
Burjassot, València, Spain*

Germán RODRIGO^a

*Institut für Theoretische Teilchenphysik, Universität Karlsruhe, 76131 Karlsruhe,
Germany*

Mikhail BILENKY^b

*Institute of Physics, AS CR, 18040 Prague 8, and Nuclear Physics Institute, AS
CR, 25068 Řež(Prague), Czech Republic*

The importance of measuring the b-quark mass at different scales is emphasized.
The recent next-to-leading order calculation of three jet heavy quark production
at LEP and its use to measure $m_b(m_Z)$ is discussed.

1 Introduction

The question of the origin of the masses of quarks and leptons is one of the unresolved puzzles in present high energy physics. To answer this question one needs to know precisely their value. However, quarks are not free and their mass has to be interpreted more like a coupling than an inertial parameter and it can run if measured at different scales. Moreover, in the standard model (SM) all fermion masses come from Yukawa couplings and those also run with the energy. To test fermion mass models one has to run masses extracted at quite different scales to the same scale and compare them with the same “ruler”. This way, for instance, one can check that in some unified models the b-quark mass and the τ -lepton mass, although different at threshold energies they could be equal at the unification scale. For instance, in fig. 1 we plot the evolution of the b-quark and the τ -lepton masses for both the standard model and the minimal supersymmetric standard model (MSSM). We see that the MSSM behaves much better than the SM since “unification” of masses could happen at much higher energies in the MSSM.

An important part of the running occurs at energies accessible in present experiments. However, the running of fermion masses, although predicted by quantum field theory, has not been tested experimentally until now. The

^aSupported in part by CSIC-Fundació Bancaixa. On leave from Departament de Física Teòrica, IFIC, Universitat de València-CSIC, 46100 Burjassot, València, Spain.

^bOn leave from JINR, 141980 Dubna, Russian Federation.




Figure 1: Running of the b-quark (dark) and τ -lepton (gray) masses for the standard model (solid) and the MSSM (dashed). b-quark mass running is given for two values of the strong coupling constant and the mass.

reason being that for energies $\sqrt{Q^2}$ much higher than the fermion mass, m_q , the mass effects become negligible since usually they are suppressed by m_q^2/Q^2 . But this argument is not always correct. In fact, in the SM, the partial decay width of the Higgs-boson into b-quarks is proportional to $m_b^2(m_H)$ because the Yukawa coupling of the Higgs-boson. In addition, it is not true¹ for jet cross sections which depend on a new variable, y_c (the jet-resolution parameter that defines the jet multiplicity) and which introduces a new scale in the analysis, $E_c = \sqrt{Q^2 y_c}$. Then, for small values of y_c there could be contributions coming like $m_q^2/E_c^2 = (m_q^2/Q^2)/y_c$ which could enhance the mass effect considerably. In addition mass effects could also be enhanced by logarithms of the mass. For instance, the ratio of the phase space for two massive quarks and a gluon to the phase space for three massless particles is $1 + 8(m_q^2/Q^2) \log(m_q/Q)$. At $Q^2 = m_Z^2$ and for the bottom quark this gives a 7% effect, for $m_b = 5$ GeV and a 3% effect for $m_b = 3$ GeV. The high precision achieved at LEP makes these effects relevant. We showed² that b-quark mass effects in three-jet production at LEP are large enough to be measured and, in fact, they were already observed in the tests of the flavour independence of $\alpha_s(m_Z)$ ^{3,4}. Therefore, these effects could be used to measure the b-quark mass at LEP.

The observable proposed as a means to extract the bottom-quark mass from LEP data was the ratio^{2,3}

$$R_3^{bd} \equiv \frac{\Gamma_{3j}^b(y_c)/\Gamma^b}{\Gamma_{3j}^d(y_c)/\Gamma^d} . \quad (1)$$

In this equation $\Gamma_{3j}^q(y_c)/\Gamma^q$ is the three-jet fraction of Z -decays and q denotes

the quark flavor. Obviously, the ratio R_3^{bd} depends on the jet-clustering algorithm⁵ used to define the jets. In this ratio and at the leading order (LO) the quark mass effects can be as large as 1% to 6%, depending on the values of the mass and the jet-resolution parameter, y_c .

Since the measurement of R_3^{bd} is done far away from the threshold of b -quark production, it can, in principle, be used to test the running of a quark mass as predicted by QCD. However, the leading order calculation does not distinguish among the different definitions of the quark mass²: perturbative pole mass, M_b , running mass at M_b -scale, or running mass at m_Z -scale. Therefore, to distinguish them it is necessary to use a complete next-to-leading order (NLO) calculation of three-jet ratios including quark masses which was not available until very recently^{6,7,8,9,10}. Here we overview the calculation⁷ used by the DELPHI Collaboration to extract the b -quark mass at the m_Z scale^{11,12,13}.

2 Jet ratios with heavy quarks at NLO

The decay width of the Z -boson into three jets with a heavy quark can be written as follows

$$\Gamma_{3j}^b = \frac{m_Z g^2 \alpha_s}{c_W^2 64 \pi^2} [g_V^2 H_V(y_c, r_b) + g_A^2 H_A(y_c, r_b)] , \quad (2)$$

where g is the SU(2) gauge coupling constant, c_W and s_W are the cosine and the sine of the weak mixing angle, $g_V = -1 + 4/3 s_W^2$ and $g_A = 1$ are the vector and axial-vector coupling of the Z -boson to the bottom quark and α_s is the strong coupling constant. Functions $H_{V(A)}(y_c, r_b)$ contain all the dependences on y_c and the quark mass, $r_b = (M_b/m_Z)^2$, for the different algorithms. These functions are computed perturbatively as an expansion in α_s and can also be expanded in r_b for $r_b \ll 1$. At the NLO we have contributions to the three-jet cross section from three- and four-parton final states. One-loop three-parton amplitudes are both infrared (IR) and ultraviolet (UV) divergent. However, the UV divergences are removed after renormalization. The four-parton transition amplitudes are also IR divergent but the sum of both, four-parton and three-parton, contributions is IR finite. We use the so-called *phase space slicing method*¹⁴ to obtain the remaining finite results, that is, the functions $H_{V(A)}$ in eq. (2) at order α_s . These results have been checked in different ways, in particular we checked that the massless result^{5,15} is nicely recovered when we take the limit $M_b \rightarrow 0$. These results have also been checked independently by two different groups^{9,10}.

Combining eq. (1), eq. (2) and using the known expression for Γ^b ^{2,16} we

write R_3^{bd} as the following expansion in α_s

$$R_3^{bd} = 1 + r_b \left(b_0 + \frac{\alpha_s}{\pi} b_1 \right) , \quad (3)$$

where the functions b_0 and b_1 are given by an average of the vector and axial-vector parts of the Z-widths, weighted by $c_V = g_V^2/(g_V^2 + g_A^2)$ and $c_A = g_A^2/(g_V^2 + g_A^2)$ respectively. It is important to note that, because of the particular normalization we have used in the definition of R_3^{bd} , most of the electroweak corrections cancel.

Although intermediate calculations have been performed using the pole mass, we can also re-express our results in terms of the running quark mass by using the known perturbative expression $M_b^2 = m_b^2(\mu)[1 + 2\alpha_s(\mu)/\pi (4/3 - \log(m_b^2/\mu^2))]$. We obtain

$$R_3^{bd} = 1 + \bar{r}_b(\mu) \left(b_0 + \frac{\alpha_s(\mu)}{\pi} \left[\bar{b}_1 - 2b_0 \log \frac{m_Z^2}{\mu^2} \right] \right) , \quad (4)$$

where $\bar{r}_b(\mu) = m_b^2(\mu)/m_Z^2$ and

$$\bar{b}_1 = b_1 + b_0 [8/3 - 2 \log(r_b)] . \quad (5)$$

$\bar{r}_b(\mu)$ can be expressed in terms of the running mass of the b -quark at $\mu = m_Z$ by using the renormalization group. At the order we are working $\bar{r}_b(\mu) = \bar{r}_b(m_Z) (\alpha_s(m_Z)/\alpha_s(\mu))^{-4\gamma_0/\beta_0}$ with $\alpha_s(\mu) = \alpha_s(m_Z)/(1 + \alpha_s(m_Z)\beta_0 t)$ and $t = \log(\mu^2/m_Z^2)/(4\pi)$, $\beta_0 = 11 - 2N_f/3$, $N_f = 5$ and $\gamma_0 = 2$.

At the perturbative level eq. (3) and eq. (4) are equivalent. However, they neglect different higher order terms and lead to different answers. The difference gives an estimate of the size of higher order corrections.

Here we present only the results for the DURHAM algorithm¹⁷, which gives smaller radiative corrections and was the one used by the DELPHI Collaboration in its analysis.

The function b_0 describes the corrections due to the quark mass at LO and has an almost negligible residual mass dependence. A fit, $b_0 = \sum_{n=0}^2 k_0^{(n)} \log^n y_c$, to the complete results for the DURHAM algorithm gives: $k_0^{(0)} = -10.521$, $k_0^{(1)} = -4.4352$, $k_0^{(2)} = -1.6629$.

The function⁷ $\bar{b}_1$ gives the NLO massive corrections to R_3^{bd} . It is important to note that $\bar{b}_1$ contains significant logarithmic corrections depending on the quark mass. We take them into account by using the form $\bar{b}_1 = k_1^{(0)} + k_1^{(1)} \log(y_c) + k_m^{(0)} \log(r_b)$ in the fit. For the DURHAM scheme we obtain: $k_1^{(0)} = 297.92$, $k_1^{(1)} = 59.358$, $k_m^{(0)} = 46.238$.




Figure 2: NLO results for R_3^{bd} (DURHAM) written in terms of the pole mass, M_b (dashed), or the running mass $m_b(m_Z)$ (solid) at LO (gray) and at the NLO (dark). The dotted line gives the NLO result in terms of the running mass at an intermediate scale $m_b(10 \text{ GeV})$. We use as a starting point $m_b(m_b) = 4.13$ to obtain both M_b and $m_b(m_Z)$.

In fig. 2 we present R_3^{bd} as a function of y_c for DURHAM. To compute it, we use the low energy measurement of the b-quark mass¹⁸, $m_b(m_b) = 4.13$. Note that choosing a low value for μ in the NLO predictions written in terms of the running mass makes it closer to the LO result written in terms of the pole mass, while choosing a large μ makes the result approach to the LO result written in terms of the running mass at the m_Z scale.

3 $m_b(m_Z)$ from LEP data

Since R_3^{bd} has been measured to good accuracy by DELPHI¹³ one can use eq. (4) and the relationship between $m_b(\mu)$ and $m_b(m_Z)$ to extract $m_b(m_Z)$. On the other hand, one could use eq. (3) to obtain directly the pole mass, M_b , and then use the relationship between the pole mass and the running mass and the renormalization group to obtain also $m_b(m_Z)$. Both results are slightly different and have a residual dependence on the scale μ . The difference gives an estimate of the errors due to the unknowledge of higher order corrections. In fig. 3 we plot, as a function of μ , the values of $m_b(m_Z)$ obtained from $R_{3exp}^{bd} = 0.971$, by using the two methods. The most conservative estimate of the theoretical error is obtained by taking the spread of the results at $\mu = m_Z$.




Figure 3: Extracted value of $m_b(m_Z)$ if $R_{3exp}^{bd} = 0.971$ as a function of the scale μ by using both the pole mass result and the running mass result.

The final result, after including also statistical errors and the errors due to the uncertainties in the hadronization corrections has been presented by DELPHI in this conference¹³. The obtained mass of the bottom-quark, $m_b(m_Z)$, measured from the three-jet decay of the Z-boson¹³ is fully compatible with the value obtained from low energy determinations¹⁸ after using the renormalization group. This provides, for the first time, a nice check of the quark mass sector of QCD in a very wide range of scales. These results can probably be improved by understanding better the scale dependence of the results, by resummation of large logs, by using other observables with a softer dependence on the scale, by reducing the hadronization uncertainties and finally by including all LEP data. We think that these studies can provide a rather precise value of $m_b(m_Z)$.

References

1. A. Ballestrero, E. Maina, and S. Moretti, *Phys. Lett. B* **294**, 425 (1992); *Nucl. Phys. B* **415**, 265 (1994).
2. M. Bilenky, G. Rodrigo, and A. Santamaria, *Nucl. Phys. B* **439**, 505 (1995).
3. DELPHI Coll., P. Abreu, et al., *Phys. Lett. B* **307**, 221 (1993)221; J. Chrin, Proc. of the 28th Rencontre de Moriond, 1993, pag. 313, ed. J.

- Trần Thanh Van; J.A. Valls, PhD. Thesis, Universitat de València, 1994.
4. L3 Coll., B. Adeva, et al. , *Phys. Lett. B* **263**, 551 (1991); OPAL Coll., R. Akers, et al., *Z. Phys. C* **65**, 31 (1995); ALEPH Coll., D. Buskulic et al., *Phys. Lett. B* **355**, 381 (1995); SLD Coll., K. Abe et al., SLAC-PUB-7573, Jan. 1997.
 5. S. Bethke, Z. Kunszt, D. E. Soper, and W. J. Stirling, *Nucl. Phys. B* **370**, 310 (1992); Z. Kunszt, P. Nason, G. Marchesini, and B. R. Webber, Proc. of the 1989 LEP Physics Workshop, Geneva, Feb 20, 1989.
 6. G. Rodrigo, *Nucl. Phys. Proc. Suppl.* **56A**, 60 (1997); PhD thesis, Universitat de València, 1996, hep-ph/9703359, ISBN: 84-370-2989-9.
 7. G. Rodrigo, A. Santamaria, and M. Bilenky, *Phys. Rev. Lett.* **79**, 193 (1997).
 8. G. Rodrigo, A. Santamaria, and M. Bilenky, hep-ph/9703360.
 9. W. Bernreuther, A. Brandenburg, and P. Uwer, *Phys. Rev. Lett.* **79**, 189 (1997).
 10. P. Nason and C. Oleari, *Phys. Lett. B* **407**, 57 (1997).
 11. J. Fuster, S. Cabrera, and S. Martí, *Nucl. Phys. Proc. Suppl.* **54A**, 39 (1997); J. Fuster, Tests of QCD at LEP, XXV international meeting on fundamental physics. Formigal – Huesca, Spain. March 3 – 8, 1997.
 12. S. Martí, J. Fuster and S. Cabrera, hep-ex/9708030; Proc. of the ICHEP'97, 19-26 August, Jerusalem, Israel, DELPHI 97-74 CONF 60.
 13. S. Cabrera, These proceedings; DELPHI Collaboration PPE-97-141, accepted in *Phys. Lett. B*.
 14. F. Aversa, M. Greco, P. Chiappetta, and J. P. Guillet, *Phys. Rev. Lett.* **65**, 401 (1990); H. Baer, J. Ohnemus, and J. F. Owens, *Phys. Rev. D* **40**, 2844 (1989); W. T. Giele and E. W. N. Glover, *Phys. Rev. D* **46**, 1980 (1992); W. T. Giele, E. W. N. Glover, and D. A. Kosower, *Nucl. Phys. B* **403**, 633 (1993); F. Gutbrod, G. Kramer, and G. Schierholz, *Z. Phys. C* **21**, 235 (1984).
 15. R. K. Ellis, D. A. Ross, and A. E. Terrano, *Nucl. Phys. B* **178**, 421 (1981).
 16. A. Djouadi, J. H. Kühn, and P. M. Zerwas, *Z. Phys. C* **46**, 411 (1990); K. G. Chetyrkin and J. H. Kühn, *Phys. Lett. B* **248**, 359 (1990).
 17. N. Brown and W. Stirling, *Z. Phys. C* **53**, 629 (1992); S. Catani, Y. L. Dokshitzer, M. Olsson, G. Turnock, and B. R. Webber, *Phys. Lett. B* **269**, 432 (1991).
 18. See for instance M. Jamin and A. Pich, *Nucl. Phys.* **B507** (1997) 334; V. Giménez, G. Martinelli, and C.T. Sachrajda, *Nucl. Phys. Proc. Suppl.* **53**, 365 (1997); *Phys. Lett. B* **393**, 124 (1997); See also S. Narison, *Phys.Lett.* **B341** (1994) 73.